

Exact time-dependent decoherence factor and its adiabatic classical limit

Jian-Qi Shen ^{*1,2}, Pan Chen ^{1,2} and Hong Mao ²

1. *Center for Optical and Electromagnetic Research, State Key Laboratory of
Modern Optical Instrumentation, College of Information Science and Engineering*

2. *Zhejiang Institute of Modern Physics and Department of Physics,
Zhejiang University, Hangzhou 310027, People's Republic of China*

(February 1, 2008)

The present paper finds the complete set of exact solutions of the general time-dependent dynamical models for quantum decoherence, by making use of the Lewis-Riesenfeld invariant theory and the invariant-related unitary transformation formulation. Based on this, the general explicit expression for the decoherence factor is then obtained and the adiabatic classical limit of an illustrative example is discussed. The result (i.e., the adiabatic classical limit) obtained in this paper is consistent with what obtained by other authors, and furthermore we obtain the more general results concerning the time-dependent non-adiabatic quantum decoherence. It is shown that the invariant theory is appropriate for treating both the time-dependent quantum decoherence and the geometric phase factor.

PACS: 03.65.Ge, 03.65.Bz

Keywords: decoherence factor; invariant theory; time-dependent decoherence

I. INTRODUCTION

Solvable models in quantum mechanics enable one to investigate quantum measurement problems very conveniently [1]. A good number of investigators have studied these useful models such as Hepp-Coleman model [2] and Cini model [3]. The exact solvability of these models often provides physicists with a clear understanding of the physical phenomena involved and yield rich physical insights [4,5]. In these works, the first important step is to obtain the exact solutions of the Schrödinger equation and the time-evolution operator, which can be applied to the calculation of the decoherence factor and study of the wavefunction collapse, etc.. Although the exact solutions and the decoherence of these models have been extensively studied by many authors, the coefficients in the Hamiltonians of all these models are merely time-independent (or partially time-dependent), to the best of our knowledge. In the present paper, we obtain the explicit time-evolution operator and the decoherence factor of the general dynamical models where the Hamiltonians are totally time-dependent.

Time-dependent system is governed by the time-dependent Schrödinger equation. The invariant theory [6] suggested by Lewis and Riesenfeld in 1969 can solve the time-dependent Schrödinger equation. In 1991, Gao *et al* proposed a generalized invariant theory [7] by introducing basic invariants, which enable one to find the complete set of commuting invariants for some time-dependent multi-dimensional systems [8–10]. We will analyze the general dynamical model in what follows and then calculate the time-dependent decoherence factor by making use of these invariant theories.

II. EXACT DECOHERENCE FACTOR IN TIME-DEPENDENT DECOHERENCE

The original Cini model for the correlation between the states of the measured system and the measuring instrument-detector is built for a two-level system interacting with the detector. Liu and Sun generalized this Cini model to an M -level system [12]. In this paper we further construct a general dynamical model for quantum decoherence between the measuring instrument-detector and the measured system where A_+ , A_- and A denote the measuring instrument-detector; and $\omega_i(t)$ represents the energy parameter of a certain state $|i\rangle$ of the multi-level measured system and $\theta_i(t)$ and $\phi_i(t)$ are coupling coefficients of interaction of the measuring instrument-detector with the measured system.

*E-mail address: jqshen@coer.zju.edu.cn

The Hamiltonian which describes the interaction between the state $|i\rangle$ of the multi-level measured system and the measuring instrument-detector is then given as follows:

$$H_i(t) = \omega_i(t) \left\{ \frac{1}{2} \sin \theta_i(t) \exp[-i\phi_i(t)] A_+ + \frac{1}{2} \sin \theta_i(t) \exp[i\phi_i(t)] A_- + \cos \theta_i(t) A \right\} \quad (2.1)$$

with A , A_+ and A_- satisfying the general commuting relations of a Lie algebra

$$[A_+, A_-] = nA, \quad [A, A_+] = mA_+, \quad [A, A_-] = -mA_-, \quad (2.2)$$

where m , $-m$ and n are structure constants of this Lie algebra. Time evolution of this dynamical model is governed by the Schrödinger equation (in the unit $\hbar = 1$)

$$i \frac{\partial |\Psi_i(t)\rangle_s}{\partial t} = H_i(t) |\Psi_i(t)\rangle_s. \quad (2.3)$$

According to the Lewis-Riesenfeld invariant theory, an operator $I(t)$ that agrees with the following invariant equation [6]

$$\frac{\partial I_i(t)}{\partial t} + \frac{1}{i} [I_i(t), H_i(t)] = 0 \quad (2.4)$$

is called an invariant whose eigenvalue is time-independent, i.e.,

$$I_i(t) |\lambda, i, t\rangle = \lambda |\lambda, i, t\rangle, \quad \frac{\partial \lambda}{\partial t} = 0. \quad (2.5)$$

It is seen from Eq. (2.4) that $I_i(t)$ is the linear combination of A_+ , A_- and A and may be generally written

$$I_i(t) = y \left\{ \frac{1}{2} \sin a_i(t) \exp[-ib_i(t)] A_+ + \frac{1}{2} \sin a_i(t) \exp[ib_i(t)] A_- \right\} + \cos a_i(t) A, \quad (2.6)$$

where the constant y will be determined below. Substitution of (2.6) into Eq. (2.4) yields

$$y \exp(-ib_i) (\dot{a}_i \cos a_i - \dot{b}_i \sin a_i) - im\omega_i [\exp(-i\phi_i) \cos a_i \sin \theta_i - y \exp(-ib_i) \sin a_i \cos \theta_i] = 0, \quad (2.7)$$

$$\dot{a}_i + \frac{ny}{2} \omega_i \sin \theta_i \sin(b_i - \phi_i) = 0. \quad (2.8)$$

where dot denotes the time derivative. The time-dependent parameters a_i and b_i are determined by these two auxiliary equations.

It is easy to verify that the particular solution $|\Psi_i(t)\rangle_s$ of the Schrödinger equation can be expressed in terms of the eigenstate $|\lambda, i, t\rangle$ of the invariant $I_i(t)$, namely,

$$|\Psi_i(t)\rangle_s = \exp\left[\frac{1}{i} \varphi_i(t)\right] |\lambda, i, t\rangle \quad (2.9)$$

with

$$\varphi_i(t) = \int_0^t \left\langle \lambda, i, t' \left| [H_i(t') - i \frac{\partial}{\partial t'}] \right| \lambda, i, t' \right\rangle dt'. \quad (2.10)$$

The physical meanings of $\int_0^t \left\langle \lambda, i, t' \left| H_i(t') \right| \lambda, i, t' \right\rangle dt'$ and $\int_0^t \left\langle \lambda, i, t' \left| -i \frac{\partial}{\partial t'} \right| \lambda, i, t' \right\rangle dt'$ are dynamical and geometric phase, respectively.

Since the expression (2.9) is merely a formal solution of the Schrödinger equation, in order to get the explicit solutions we make use of the invariant-related unitary transformation formulation [7] which enables one to obtain the complete set of exact solutions of the time-dependent Schrödinger equation (2.3). In accordance with the invariant-related unitary transformation method, the time-dependent unitary transformation operator is often of the form

$$V_i(t) = \exp[\beta_i(t) A_+ - \beta_i^*(t) A_-] \quad (2.11)$$

with $\beta_i(t) = -\frac{a_i(t)}{2}x \exp[-ib_i(t)]$, $\beta_i^*(t) = -\frac{a_i(t)}{2}x \exp[ib_i(t)]$. By making use of the Glauber formula, lengthy calculation yields

$$\begin{aligned}
I_{iV} &= V_i^\dagger(t)I_i(t)V_i(t) = \left\{ \frac{y}{2} \exp(-ib_i) \sin a_i \cos\left[\left(\frac{mn}{2}\right)^{\frac{1}{2}} a_i x\right] \right. \\
&\quad - \frac{\left(\frac{mn}{2}\right)^{\frac{1}{2}}}{n} \exp(-ib_i) \cos a_i \sin\left[\left(\frac{mn}{2}\right)^{\frac{1}{2}} a_i x\right] \Big\} A_+ \\
&\quad + \left\{ \frac{y}{2} \exp(ib_i) \sin a_i \cos\left[\left(\frac{mn}{2}\right)^{\frac{1}{2}} a_i x\right] \right. \\
&\quad - \frac{\left(\frac{mn}{2}\right)^{\frac{1}{2}}}{n} \exp(ib_i) \cos a_i \sin\left[\left(\frac{mn}{2}\right)^{\frac{1}{2}} a_i x\right] \Big\} A_- \\
&\quad + \left\{ \cos a_i \cos\left[\left(\frac{mn}{2}\right)^{\frac{1}{2}} a_i x\right] + \frac{\left(\frac{mn}{2}\right)^{\frac{1}{2}}}{m} y \sin a_i \sin\left[\left(\frac{mn}{2}\right)^{\frac{1}{2}} a_i x\right] \right\} A.
\end{aligned} \tag{2.12}$$

It can be easily seen that when

$$y = \frac{m}{\left(\frac{mn}{2}\right)^{\frac{1}{2}}}, \quad x = \frac{1}{\left(\frac{mn}{2}\right)^{\frac{1}{2}}}, \tag{2.13}$$

one may derive that $I_{iV} = A$ which is time-independent. Thus the eigenvalue equation of the time-independent invariant I_{iV} may be written in the form

$$I_{iV} |\lambda\rangle = \lambda |\lambda\rangle, \quad |\lambda\rangle = V_i^\dagger(t) |\lambda, i, t\rangle. \tag{2.14}$$

In the meanwhile, by the aid of Baker-Campbell-Hausdorff formula [13], one can arrive at

$$\begin{aligned}
V_i^\dagger(t)H_i(t)V_i(t) &= \left\{ \omega_i \left[\frac{1}{2} \sin \theta_i \exp(-i\phi_i) - \frac{\sqrt{\frac{mn}{2}}}{n} \exp(-ib_i) \cos \theta_i \sin\left(\sqrt{\frac{mn}{2}} a_i x\right) \right] \right. \\
&\quad + \frac{1}{2} \omega_i \exp(-ib_i) \sin \theta_i \cos(b_i - \phi_i) [\cos(\sqrt{\frac{mn}{2}} a_i x) - 1] \Big\} A_+ \\
&\quad + \left\{ \omega_i \left[\frac{1}{2} \sin \theta_i \exp(i\phi_i) - \frac{\sqrt{\frac{mn}{2}}}{n} \exp(ib_i) \cos \theta_i \sin\left(\sqrt{\frac{mn}{2}} a_i x\right) \right] \right. \\
&\quad + \frac{1}{2} \omega_i \exp(ib_i) \sin \theta_i \cos(b_i - \phi_i) [\cos(\sqrt{\frac{mn}{2}} a_i x) - 1] \Big\} A_- \\
&\quad + \omega_i [\cos \theta_i \cos(\sqrt{\frac{mn}{2}} a_i x) + \frac{\sqrt{\frac{mn}{2}}}{m} \sin \theta_i \cos(b_i - \phi_i) \sin(\sqrt{\frac{mn}{2}} a_i x)] A,
\end{aligned} \tag{2.15}$$

which is related to the dynamical phase; and

$$\begin{aligned}
V_i^\dagger(t)i\frac{\partial V_i(t)}{\partial t} &= \left[-\frac{i}{2} \dot{a}_i x \exp(-ib_i) - \frac{1}{2} \frac{1}{\sqrt{\frac{mn}{2}}} \dot{b}_i \exp(-ib_i) \sin\left(\sqrt{\frac{mn}{2}} a_i x\right) \right] A_+ \\
&\quad + \left[\frac{i}{2} \dot{a}_i x \exp(ib_i) - \frac{1}{2} \frac{1}{\sqrt{\frac{mn}{2}}} \dot{b}_i \exp(ib_i) \sin\left(\sqrt{\frac{mn}{2}} a_i x\right) \right] A_- \\
&\quad - \frac{\dot{b}_i}{m} [1 - \cos(\sqrt{\frac{mn}{2}} a_i x)] A,
\end{aligned} \tag{2.16}$$

which is related to the geometric phase. It follows from Eq. (2.15) and Eq. (2.16) that, under the transformation $V(t)$, the Hamiltonian $H(t)$ can be changed into

$$\begin{aligned}
H_{iV}(t) &= V_i^\dagger(t)H_i(t)V_i(t) - V_i^\dagger(t)i\frac{\partial V_i(t)}{\partial t} \\
&= \left\{ \omega_i [\cos a_i \cos \theta_i + \frac{\left(\frac{mn}{2}\right)^{\frac{1}{2}}}{m} \sin a_i \sin \theta_i \cos(b_i - \phi_i)] \right. \\
&\quad \left. + \frac{\dot{b}_i}{m} (1 - \cos a_i) \right\} A
\end{aligned} \tag{2.17}$$

Hence, with the help of Eq. (2.9) and Eq. (2.14), the particular solution of the Schrödinger equation is obtained

$$|\Psi_i(t)\rangle_s = \exp\left[\frac{1}{i}\varphi_i(t)\right]V_i(t)|\lambda, i\rangle \quad (2.18)$$

with the phase

$$\begin{aligned} \varphi_i(t) &= \int_0^t \langle \lambda | [V_i^\dagger(t')H_i(t')V_i(t') - V_i^\dagger(t')i\frac{\partial}{\partial t'}V_i(t')] | \lambda \rangle dt' \\ &= \varphi_{id}(t) + \varphi_{ig}(t) \\ &= \lambda \int_0^t \left\{ \omega_i [\cos a_i \cos \theta_i + \frac{(\frac{mn}{2})^{\frac{1}{2}}}{m} \sin a_i \sin \theta_i \cos(b_i - \phi_i)] \right. \\ &\quad \left. + \frac{\dot{b}_i}{m}(1 - \cos a_i) \right\} dt', \end{aligned} \quad (2.19)$$

where the dynamical phase is $\varphi_{id}(t) = \lambda \int_0^t \omega_i [\cos a_i \cos \theta_i + \frac{(\frac{mn}{2})^{\frac{1}{2}}}{m} \sin a_i \sin \theta_i \cos(b_i - \phi_i)] dt'$ and the geometric phase is $\varphi_{ig}(t) = \lambda \int_0^t \frac{\dot{b}_i}{m}(1 - \cos a_i) dt'$. It is seen that the former phase is related to the dynamical parameters of the Hamiltonian such as $\omega_i, \cos \theta_i, \sin \theta_i$, etc., whereas the latter is not immediately related to these parameters. If the parameter a_i is taken to be time-independent, then we arrive at

$$\varphi_{ig}(T) = \lambda \int_0^T \frac{\dot{b}_i}{m}(1 - \cos a_i) dt' = \frac{\lambda}{m} [2\pi(1 - \cos a_i)], \quad (2.20)$$

where $2\pi(1 - \cos a_i)$ is an expression for the solid angle over the parameter space of the invariant. This fact shows the global or topological meanings of the geometric phase $\varphi_{ig}(t)$ [11]. The expression (2.18) is a particular exact solution corresponding to λ and the general solutions of the time-dependent Schrödinger equation are easily obtained by using the linear combinations of all these particular solutions.

Since we have exact solutions of the general time-dependent model, we can obtain the exact expression for the time-dependent decoherence factor that is given

$$F_{i,j}(t) = \langle \lambda | V_i^\dagger(t)V_j(t) | \lambda \rangle. \quad (2.21)$$

Further calculation yields

$$F_{i,j}(t) = \exp\left[\frac{n\lambda}{2}(\beta_i\beta_j^* - \beta_i^*\beta_j)\right] \langle \lambda | \exp[(\beta_j - \beta_i)A_+ - (\beta_j^* - \beta_i^*)A_-] | \lambda \rangle, \quad (2.22)$$

which is the general expression for the decoherence factor of the time-dependent dynamical model (2.1). Although the expression (2.5) is somewhat complicated, it is just the explicit expression that does not contain the chronological product.

III. AN ILLUSTRATIVE EXAMPLE

To show that (2.5) descends to the result familiar to us in the time-independent (or partially time-dependent) dynamical model of decoherence, we take into consideration the adiabatic classical limit of a special dynamical model. Liu and Sun generalized the original Cini model to an M -level system [12]. The Hamiltonian of this generalized model is written

$$H = H_S + H_D + H_I, \quad (3.1)$$

where H_S is the model Hamiltonian of the measured system S with M levels and H_D is the free Hamiltonian of the two-boson-state detector D. They are generally of the forms

$$H_S = \sum_{k=1}^M E_k |\Phi_k\rangle \langle \Phi_k|, \quad H_D = \omega_1 a_1^\dagger a_1 + \omega_2 a_2^\dagger a_2 \quad (3.2)$$

with the creation and annihilation operators $a_i^\dagger$, a_i satisfying the following commuting relations

$$[a_i, a_j^\dagger] = \delta_{ij}, \quad [a_i, a_j] = [a_i^\dagger, a_j^\dagger] = 0. \quad (3.3)$$

The interaction Hamiltonian H_I is given by

$$\begin{aligned} H_I &= \sum_n |\Phi_n\rangle \langle \Phi_n| (g_n a_1^\dagger a_2 + g_n^* a_2^\dagger a_1) \\ &= \sum_n |\Phi_n\rangle \langle \Phi_n| (g_n J_+ + g_n^* J_-) \end{aligned} \quad (3.4)$$

with $J_+ = a_1^\dagger a_2$, $J_- = a_2^\dagger a_1$, $J_3 = \frac{1}{2}(a_1^\dagger a_1 - a_2^\dagger a_2)$ satisfying the commuting relations $[J_+, J_-] = 2J_3$, $[J_3, J_\pm] = \pm J_\pm$.

It can be seen from the form of the Hamiltonian that both $|\Phi_k\rangle \langle \Phi_k|$ and $N = \frac{a_1^\dagger a_1 + a_2^\dagger a_2}{2}$ commute with H , namely, $[|\Phi_k\rangle \langle \Phi_k|, H] = [N, H] = 0$. Hence, a generalized quasialgebra which enables one to obtain the complete set of exact solutions of the Schrödinger equation can be found by working in a sub-Hilbert-space corresponding to the particular eigenvalues of both $|\Phi_k\rangle \langle \Phi_k|$ and N , and then the Hamiltonian can be rewritten in this sub-Hilbert-space

$$H_{n,k}(t) = E_k + g_k J_+ + g_k^* J_- + (\omega_1 - \omega_2) J_3 + n(\omega_1 + \omega_2) \quad (3.5)$$

with n being the eigenvalue of N and satisfying

$$N |n_1, n_2\rangle = n |n_1, n_2\rangle, \quad n = \frac{1}{2}(n_1 + n_2). \quad (3.6)$$

Thus in the sub-Hilbert-space we write the Schrödinger equation in the form

$$H_{n,k}(t) |\Psi_{n,k}(t)\rangle_s = i \frac{\partial}{\partial t} |\Psi_{n,k}(t)\rangle_s, \quad (3.7)$$

and $|\Psi(t)\rangle_s$ can be obtained from

$$|\Psi(t)\rangle_s = \sum_n \prod_k c_{n,k} |\Psi_{n,k}(t)\rangle_s |\Phi_k\rangle \quad (3.8)$$

where $c_{n,k}$ is time-independent and determined by the initial conditions.

For the case of this time-dependent Cini model, we set $A_+ = J_+$, $A_- = J_-$, $C = J_z$ with $J_\pm = J_1 \pm iJ_2$. In the adiabatic limit, it follows from the auxiliary equations (2.8) that

$$a_i = \theta_i, \quad b_i = \varphi_i; \quad a_j = \theta_j, \quad b_j = \varphi_j. \quad (3.9)$$

Since

$$\beta_i = -\frac{a_i}{2} \exp[-ib_i], \quad \beta_j = -\frac{a_j}{2} \exp[-ib_j], \quad (3.10)$$

we let both b_i and b_j vanish for the convenience, which leads to $\dot{\theta}_i = 0$ and $\dot{\theta}_j = 0$ in terms of the auxiliary equations (2.8), and then $\beta_i = -\frac{a_i}{2}$, $\beta_j = -\frac{a_j}{2}$. It is therefore easily obtained that

$$\begin{aligned} F_{i,j}(t) &= \exp\left[\frac{\lambda}{2}(\beta_i \beta_j^* - \beta_i^* \beta_j)\right] \langle j, m | \exp[(\beta_j - \beta_i)J_+ - (\beta_j^* - \beta_i^*)J_-] | j, m \rangle \\ &= \langle j, m | \exp\{[(\beta_j - \beta_i) - (\beta_j^* - \beta_i^*)]J_1 \\ &\quad + i[(\beta_j - \beta_i) + (\beta_j^* - \beta_i^*)]J_2\} | j, m \rangle \\ &= \langle j, m | \exp[i(a_i - a_j)J_2] | j, m \rangle, \end{aligned} \quad (3.11)$$

where j and m satisfy

$$J_3 |j, m\rangle = m |j, m\rangle, \quad J^2 |j, m\rangle = j(j+1) |j, m\rangle. \quad (3.12)$$

If when $t = 0$, then the state of the measuring instrument-detector is $|j, j\rangle$, and the decoherence factor is therefore

$$\begin{aligned}
F_{i,j}(t) &= \langle j, j | \exp[i(a_i - a_j)J_2] | j, j \rangle \\
&= [\cos(\frac{a_i - a_j}{2})]^{2j},
\end{aligned} \tag{3.13}$$

which is consistent with that obtained by Sun *et al* [14]. For the case of classical limit where $j \rightarrow \infty$ and $\frac{a_i - a_j}{2} \neq n\pi$ ($n = 0, \pm 1, \pm 2, \dots$), $F_{k,l}(t) \rightarrow 0$, which means the wavefunction collapse occurs under the classical limit.

Since we exactly solved the Schrödinger equation governing the time-dependent quantum decoherence, and the result (i.e., the adiabatic classical limit) obtained here is consistent with what obtained in previous references, we hold that the more general results presented in this paper is useful to treat the time-dependent non-adiabatic quantum decoherence.

IV. CONCLUDING REMARKS

The present paper obtains exact solutions and decoherence factor of the general time-dependent dynamical model for quantum decoherence by working in the sub-Hilbert space corresponding to the eigenvalue of two invariants and by making use of the invariant-related unitary transformation method. The invariant-related unitary transformation formulation is an effective method for treating time-dependent problems [15–17]. This formulation replaces eigenstates of the time-dependent invariants with those of the time-independent invariants through the unitary transformation. It uses the invariant-related unitary transformation and obtains the explicit expression for the time-evolution operator, instead of the formal solution that is related to the chronological product. In view of what has been discussed above, it can be seen that the invariant theory is appropriate to treat the time-dependent quantum decoherence. Apparently, the results presented in the present paper is easy to generalize to the time-dependent Hepp-Coleman model. Since the geometric phase factor appears in time-dependent systems, it is interesting to investigate the geometric phase in the time-dependent quantum decoherence by using the formulation in this paper.

Acknowledgment This project was supported by the National Natural Science Foundation of China under the project No. 30000034.

-
- [1] H. Nakazato, S. Pascazio, Phys. Rev. Lett. 70 (1993) 1.
 - [2] M. Namiki, S. Pascazio, Phys. Rev. A 44 (1991) 39.
 - [3] M. Cini, Nuovo Cimento. B 73 (1983) 27.
 - [4] M. Namiki, S. Pascazio, Found. Phys.Lett. 4 (1991) 203.
 - [5] H. Nakazato, S. Pascazio. Phys. Rev. A 57 (1998) 753.
 - [6] H. R. Lewis, W. B. Riesenfeld, J. Math. Phys. 10 (1969) 1458.
 - [7] X. C. Gao, J. B. Xu, T. Z. Qian, Phys. Rev. A 45 (1992) 4355.
 - [8] X. C. Gao, J. Fu, J. Q. Shen, Eur. Phys. J. C 13 (2000) 527.
 - [9] J. Q. Shen, H. Y. Zhu, J. Li, Acta Phys. Sin. 50 (2001) 1884.
 - [10] S. P. Kim, A. E. Santana, F. C. Khanna, Phys. Lett. A 272 (2000) 46.
 - [11] M. V. Berry, Proc. R. Soc. Lond. A 392 (1984) 45.
 - [12] X. J. Liu, C. P. Sun, Phys. Lett. A 198 (1995) 371.
 - [13] J. Wei, E. Norman, J. Math. Phys.(N.Y) 4 (1963) 575.
 - [14] J. Y. Zeng, S. Y. Pei, New Advance in Quantum Mechanics, Publishing House of Pecking University, Pecking, 2000 (Chapter 3).
 - [15] J. Fu, X. C. Gao, J. B. Xu, X. B. Zou, Can. J. Phys. 77 (1999) 1.
 - [16] J. Q. Shen, H. Y. Zhu, S. L. Shi, J. Li, Phys. Scr. 65 (2002) 465.
 - [17] J. Q. Shen, H. Y. Zhu, S. L. Shi, J. Li, J. Phys. Soc. Jpn. 71 (2001) 1440.